

# Direction fields, autonomous ODEs, and the phase line Lecture 7e – 2021-07-07

MAT A35 – Summer 2021 – UTSC


Prof. Yun William Yu

# 1<sup>st</sup>-order ODEs and slopes of solutions

- We can write a 1<sup>st</sup>-order ODE as  $y' = f(x, y)$
- Recall that the derivative can be thought of as the slope of a solution.

# Direction field

- A direction field graphs out the slopes of all solutions going through a point.
- We can visualize different solutions by drawing trajectory curves that are always tangent to the direction field.



# Autonomous ODEs


- Recall that an autonomous ODE is one that does not have an explicit dependence on the independent variable (e.g. time).
- A first-order autonomous ODE can be rewritten in the form:

$$y' = f(y)$$


# Direction fields of autonomous ODEs

- Notice that if  $y' = f(y)$ , then the slope has no  $x$ -dependence.

$$y' = \sin y$$



$$y' = y^2$$




# Equilibrium values

- An equilibrium value of the autonomous ODE  $y' = f(y)$  is a constant solution  $y = c$ .
- We can solve for equilibrium values by setting  $y' = 0$ .

# Try it out: find the equilibrium values

- $y' = y^2 - y - 6$



- A: -2
- B: 3
- C: All of the above
- D: ???
- E: None of the above

# Stability of equilibrium values


- Consider an equilibrium value  $c$  of  $y' = f(y)$ , and an initial value  $y(0) = y_0$ , where  $y_0 \approx c$ , but  $y_0 \neq c$ . Then  $c$  is
  - Unstable if  $y$  diverges from  $c$  as time  $x \rightarrow \infty$
  - Asymptotically stable if  $y(x) \rightarrow c$  as  $x \rightarrow \infty$
  - Semi-stable if as  $x \rightarrow \infty$ ,  $y(x)$  goes to  $c$  on one side, but diverges on the other side.



# Determining stability using sign of $y'$

- $y' = 0$  at equilibrium.
- $y' > 0$  implies  $y(x)$  gets bigger
- $y' < 0$  implies  $y(x)$  gets smaller


$$y' = y^2 - y - 6$$



# Phase line

- Draw arrows along the  $y$ -axis depending on if  $z = f(y) = y'$  is positive or negative.

$$y' = y^4 - 4y^2$$




# Phase line stability of $y' = f(y)$

- If  $z = f(y)$  crosses the  $y$ -axis at  $f(c) = 0$  going upward, then  $c$  is unstable.
- If  $z = f(y)$  crosses the  $y$ -axis at  $f(c) = 0$  going downward, then  $c$  is asymptotically stable.
- If  $z = f(y)$  touches the  $y$ -axis at  $f(c) = 0$ , but remains on the same side of the  $y$ -axis, then  $c$  is semi-stable.

# Try it out

- $y' = y^5 - 3y^4 - 4y^3 + 12y^2$
- Classify the stability of the following equilibria:
- $y = -2$
- $y = 0$
- $y = 2$
- $y = 3$




- A: Asymptotically stable
- B: Unstable
- C: Semi-stable
- D: ???
- E: None of the above

# Derivative test for stability

- Let  $y' = f(y)$  have an equilibrium at  $f(c) = 0$ .
- If  $\frac{df}{dy}(c) > 0$ , then  $y = c$  is unstable.
- If  $\frac{df}{dy}(c) < 0$ , then  $y = c$  is asymptotically stable.
- If  $\frac{df}{dy}(c) = 0$  and  $c$  is a local extremum (max or min) of  $f(y)$ , then  $y = c$  is semi-stable

# Example

- $y' = y^5 - 3y^4 - 4y^3 + 12y^2$  has equilibria -2, 0, 2, 3



# Try it out


- Fat crystallization: Let  $y(x)$  be the proportion of crystallizable milk fat in a sample after  $x$  hours, satisfying
$$y' = 8(y^5 - y)$$
- If you start with half of the fat as crystallizable, how much fat is crystallizable as time goes to  $\infty$ ?



[https://commons.wikimedia.org/wiki/File:Glass\\_of\\_milk.jpg](https://commons.wikimedia.org/wiki/File:Glass_of_milk.jpg)

- A: All of the fat
- B: Half of the fat
- C: None of the fat
- D: ???
- E: None of the above

# Logistic Growth Model



- Previously, we saw exponential growth  $y' = ky$ ,  $y(0) = y_0$ , which had a solution  $y(x) = y_0 e^{kx}$ .
- In practice, this is unrealistic. For example, bacteria in a petri dish will initially grow almost exponentially, but then they'll use up all the available media.
- A better model is the logistic model,  $y' = ky \left(1 - \frac{y}{L}\right)$ , where the parameter  $L$  is the carrying capacity of the environment, and  $k > 0$  is still the growth rate.



# Stability of equilibria of logistic model

- $y' = ky \left(1 - \frac{y}{L}\right)$

# Logistic model behavior

- $y' = ky \left(1 - \frac{y}{L}\right)$
- $y < 0$  is not physical, since we cannot have negative bacteria.
- When  $0 < y < L$ ,  $y' > 0$ , so the number of bacteria increase, up to  $L$ .
- When  $y > L$ ,  $y' < 0$ , so the number of bacteria decrease, down to  $L$ .
- On large time-scales, we therefore have  $L$  bacteria.