

## Quiz 1 - Thursday - Solutions

### Problem 1 (40pts)

Solve each of the following problems. If there are multiple potential answers, give your response in the most general form possible.

- $\int 2dx = 2x + C$
- $\int x^{-3}dx = -\frac{1}{2}x^{-2} + C$
- $\int \frac{2}{2x-1}dx = \ln|2x-1| + C$
- $\int \frac{1}{2}\cos(x+1)dx = \frac{1}{2}\sin(x+1) + C$
- $\int (x^2 - 4x + 4)dx = \frac{1}{3}x^3 - 2x^2 + 4x + C$
- $\int \frac{1}{(y+1)(2y+1)}dy = \ln|2y+1| - \ln|y+1| + C$
- $\int (e^{-2u} + u^3)du = -\frac{1}{2}e^{-2u} + \frac{1}{4}u^4 + C$
- $\int \left[ \frac{d}{dx}((\cos(5 - \sin x))e^{x^2} - 4) \right] dx = (\cos(5 - \sin x))e^{x^2} + C$

### Problem 2 (30pts)

Solve each of the following definite integrals. You should simplify as much as possible without a calculator, but may leave answers in terms of  $e$ ,  $\ln$ ,  $\sin$ ,  $\sqrt{\dots}$ , etc.

Each problem is worth 15 points.

- $\int_0^{\pi/2} (1 + \sin x)^2 \cos x dx$

Take off additional points for arithmetic errors, but don't penalize twice. i.e. if they made a silly error early on, but got the rest of the problem right, give them most of the later credit.

Option 1 (change the limits)

- u-substitution  $u = 1 + \sin x$ ,  $du = \cos x dx$
- changing the limits to  $u = 1, 2$ :  $\int_1^2 u^2 du$
- solving to get  $\frac{7}{3}$

Option 2 (solve the antiderivative first and change back to x's)

- u-substitution  $u = 1 + \sin x$ ,  $du = \cos x dx$
- solving to get  $\frac{1}{3}u^3 = \frac{1}{3}(1 + \sin x)^3$
- plugging in the original limits to get  $\frac{7}{3}$ .

- $\int_0^{\pi} xe^{-4x} dx + \int_{\pi}^{\infty} xe^{-4x} dx$


- recognizing that the integrals can be combined to just  $[0, \infty]$
- integration by parts using  $u = x$ ,  $du = dx$  and  $dv = e^{-4x}$ ,  $v = -\frac{1}{4}e^{-4x}$ . Should get  $-\frac{x}{4}e^{-4x} - \frac{1}{16}e^{-4x}$ .
- solving the improper integral by "plugging-in" 0 and  $\infty$  to get  $\frac{1}{16}$



### Problem 3 (15pts)

Every day, the Gibson coal-fired power plant in Owensville, Indiana, USA produces about 50 kilotons of  $\text{CO}_2$  into the atmosphere.

Suppose you set up a post-combustion carbon capture system set up. Starting from midnight, the instantaneous rate of carbon capture is  $f(t) = te^{-t} + 2 + e^{t-24}$  kilotons/hour of  $\text{CO}_2$ .



After carbon capture, how much  $\text{CO}_2$  would be released by the power plant into the air each day? You may approximate  $e^{-24} \approx 0$ .

- recognizing that total carbon capture per day is  $\int_0^{24} f(t)dt$ .
- $\int_0^{24} 2dt = 48$ .
- $\int_0^{24} e^{t-24}dt = 1 - \frac{1}{e^{24}} \approx 1$
- $\int_0^{24} te^{-t}dt = 1 - \frac{25}{e^{24}} \approx 1$
- $50 - (48 + 1 + 1) = 0$ . So just about all of the  $\text{CO}_2$  is captured.



### Problem 4 (15pts)

You are King Arthur's court wizard. Your job is to determine the average airspeed of an unladen European swallow, but you only have one swallow. You decide to send it on a 2-hour round-trip journey, and you measure its speed twice an hour, for a total of 5 measurements (including the 0th-hour measurement).

- $v(0) = 22$  miles per hour
- $v(0.5) = 25$  miles per hour
- $v(1) = 18$  miles per hour
- $v(1.5) = 27$  miles per hour
- $v(2.0) = 22$  miles per hour

1. Approximate the distance traveled using rectangular Riemann sums. Use this to compute the average speed.

The student may either use right- or left-rectangular Riemann sums. Note that getting average speed by just taking the 5 speed measurements and averaging them is INCORRECT for the purposes of this problem.

**Right Riemann sums:**

- total distance =  $(0.5 \text{ hour})(25+18+27+22 \text{ mph}) = 46$  miles.
- average speed =  $46 \text{ miles} / 2 \text{ hours} = 23 \text{ mph}$

**Left Riemann sums:**

- total distance =  $(0.5 \text{ hour})(22+25+18+27 \text{ mph}) = 46$  miles.
- average speed =  $46 \text{ miles} / 2 \text{ hours} = 23 \text{ mph}$

2. Approximate the distance traveled using the trapezoid rule. Use this to compute the average speed.

Trapezoid rule requires adding all datapoints times two, except for the first and last ones. Then divide by two.

- $(0.5 \text{ hours})[22 + 2(25+18+27) + 22 \text{ mph}]/2 = 92/2 = 46 \text{ miles}$  (note that remembering the multiplication/division by 2 for the trapezoid rule is worth 3 pts. i.e. they have to show that they actually remembered the trapezoid rule)
- average speed = 46 miles / 2 hours = 23 mph.

In this particular problem, it turns out that you get the same answer if you use Riemann sums or the trapezoid rule. Thus, the student must demonstrate that they actually applied the appropriate rule to get credit.