

Phase portraits

Lecture 10c: 2023-03-23

MAT A35 – Winter 2023 – UTSC

Prof. Yun William Yu

System of two 1st-order ODEs

- $$\begin{cases} \dot{x} = x + y - \sin t \\ \dot{y} = x^2 + y^2 - \ln t \end{cases}$$
(nonautonomous)
- How many dimensions do nonautonomous systems need to draw direction fields?
- $$\begin{cases} \dot{x} = x + y \\ \dot{y} = x^2 + y^2 \end{cases}$$
(autonomous system)
- How many dimensions do autonomous systems need to draw direction fields?
- How many dimensions do autonomous systems need to draw phase “lines”?

A: 1

B: 2


C: 3

D: 4

E: None of the above

Plotting vector fields and trajectories

- Consider $\begin{cases} \dot{x} = f(x, y) \\ \dot{y} = g(x, y) \end{cases}$
- The system associates a direction and a magnitude for every point in \mathbb{R}^2 , telling you what direction trajectories go.
- WolframAlpha: “vector field {f(x,y), g(x,y)}”
- Ex: “vector field {y, -x}” or “integral curves {y, -x}”
- Specify limits by adding “x=-3..3, y=-3..3” after




Try it out


- Which of the following is the integral curves for the system, plotted for x and y both between -5 and 5?

$$\begin{aligned}\dot{x} &= 4x - y - \left(x + \frac{3}{2}y\right)(x^2 + y^2) \\ \dot{y} &= x + 4y + \left(\frac{3}{2}x - y\right)(x^2 + y^2)\end{aligned}$$


A:




B:



C:



D:





https://www.wolframalpha.com/input/?i=integral+curves+%7B4*x-y-%28x%2B1.5y%29*%28x%5E2%2By%5E2%29%2C+x%2B4*y%2B%281.5x-y%29*%28x%5E2%2By%5E2%29%7D%2C+x%3D-5..5%2C+y%3D-5..5

Phase plane analysis

- Consider the autonomous homogeneous 2D linear system with constant coefficients

$$\begin{bmatrix} \dot{x} \\ \dot{y} \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}, A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

- Also, notice that the origin is always an equilibrium for a linear system.



Using the eigendecomposition

- If (λ_1, v_1) and (λ_2, v_2) is an eigendecomposition of A , then the general solution describing any trajectory is

$$c_1 e^{\lambda_1 t} v_1 + c_2 e^{\lambda_2 t} v_2$$

- We can qualitatively analyze the behavior of the system by looking at the eigenvalues and eigenvectors.
- Consider $f(t) = ce^{\lambda t}$.
- If $\lambda > 0$, we get exponential growth away from 0.
- If $\lambda < 0$, we get exponential decay towards 0.

Sign and stability

- If eigenvalues are positive (or have a positive real part), then trajectories go away from the origin. (unstable node)
- If eigenvalues are negative (or have a negative real part), then trajectories go towards the origin. (asymptotically stable node)
- If eigenvalues have opposite signs, then we have a saddle point, as trajectories come in along one eigenvector, and leave along the other. (unstable, saddle point)

Complex eigenvalues

- Recall complex eigenvalues come in pairs $\lambda_{1,2} = a \pm bi$.


- Solutions look like

$$z = c_1 v_1 e^{at} \cos bt + c_2 v_2 e^{at} \sin bt$$

- The sign of the real part a determines if the trajectories go inward (stable) or outward (unstable).
- The imaginary term means that the trajectories have a rotational component; i.e. might spiral in or out, or form a circle.

(Im)proper nodes

- Sometimes, if $\lambda_1 = \lambda_2$, there is only one eigenvector. Then we have an *improper* node that's hard to draw.
 - Sign still determines stable vs unstable.
- If $\lambda_1 = \lambda_2$ and we have two eigenvectors, then we have a *proper* node, which looks like a star.



Summarizing everything

- $\begin{bmatrix} \dot{x} \\ \dot{y} \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}, A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$
- The origin (0,0) is always an equilibrium point.
- We can understand the behavior around the origin by looking at the eigenvalues of A .
- Positive real parts mean that the trajectories go outward.
- Negative real parts mean that the trajectories go inward.
- Opposite sign eigenvalues mean you have a saddle point.
- Nonzero imaginary components mean that trajectories spiral.

Try it out

- $\lambda_1 = 4, \lambda_2 = -2$
- $\lambda_1 = -3, \lambda_2 = -1$
- $\lambda_1 = 2, \lambda_2 = 3$
- $\lambda_1 = 3, \lambda_2 = 3$
- $\lambda_1 = 3 + 2i, \lambda_2 = 3 - 2i$
- $\lambda_{1,2} = -1 \pm 2i$
- $\lambda_{1,2} = \pm 4i$

- A: Asymptotically Stable
- B: Stable
- C: Unstable
- D: ???
- E: None of the above

Special note: weird stuff can happen when $\lambda = 0$, which we won't deal with.

- A: Node (incl. (im)proper)
- B: Saddle Point
- C: Spiral
- D: Center
- E: None of the above

Example

- Classify the behavior around the origin of

$$\begin{bmatrix} \dot{x} \\ \dot{y} \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

Example

- Classify the behavior around the origin of

$$\begin{bmatrix} \dot{x} \\ \dot{y} \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

Try it out

- Classify the behavior around the origin of

$$\begin{bmatrix} \dot{x} \\ \dot{y} \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ -3 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$$

A: Asymptotically Stable
B: Stable
C: Unstable
D: ???
E: None of the above

A: Node (incl. (im)proper)
B: Saddle Point
C: Spiral
D: Center
E: None of the above