


Multilinear and Nonlinear Regression Lecture 6c: 2023-02-16

MAT A35 – Winter 2023 – UTSC

Prof. Yun William Yu


Single variable linear regression

- Given samples of the dependent variable y_1, \dots, y_n at values of the independent variable x_1, \dots, x_n , we want to find the linear model $f(x) = mx + b$ such that $y_i \approx f(x_i)$, the “best-fit” line.




Two-variable linear regression


- What if we have multiple independent variables?
- Suppose we are measuring the water temperature in Lake Ontario, and want to know how the temperature varies as a function of location



3D Scatter Plot of temperatures



Best-fit plane





Two-variable linear regression

- Let x and y be the independent variables. Let z be the dependent variable. Given samples z_1, \dots, z_n at values $(x_1, y_1), \dots, (x_n, y_n)$, we want the linear model

$$f(x, y) = m_1x + m_2y + b$$

such that $z_i \approx f(x_i, y_i)$, the “best-fit” plane.



Multilinear regression

- One independent variable, one dependent variable
- Two independent variables, one dependent variable
- Many independent variables, one dependent variable
- Can also have many independent variables, many dependent...

Try it out

- You are measuring the temperature of Lake Ontario as a function of location. You get the following data:


Longitude	Latitude	Temperature
76.5 W	43.5 N	12.2
76.5 W	43.9 N	12.1
77.0 W	43.6 N	11.6
77.0 W	43.8 N	11.5
78.0 W	43.3 N	13.7
78.0 W	43.7 N	13.1
79.5 W	43.8 N	12.3
79.5 W	43.9 N	12.1

- A: 12.06
- B: 12.35
- C: 12.54
- D: 12.89
- E: None of the above

- The GPS coordinates for the lake near Toronto are 43.6 N, 79.3 W. What do you predict the lake water temperature to be near Toronto?

Polynomial regression

- What if our data doesn't look linear?



Different types of regression

- Linear regression:

$$f(x) = mx + b$$

- Quadratic regression:

$$f(x) = m_2x^2 + m_1x + b$$

- Cubic regression:

$$f(x) = m_3x^3 + m_2x^2 + m_1x + b$$

- Polynomial regression of degree n :

$$f(x) = b + \sum_{i=1}^n m_i x^i$$

- Exponential regression:

$$f(x) = b + m_0 e^{m_1 x}$$

Convert nonlinear to multilinear

Intuition guess

- Linear vs. Quadratic vs Cubic: which model will have smaller Mean Square Error for the following data:

A: Linear
B: Quadratic
C: Cubic
D: Same error for All
E: None of the above

Be careful about too many parameters

- The more parameters you have (e.g. in a polynomial regression), the better your mean squared error will be.
- However, sometimes, you will overfit to the data.
- John von Neumann: “with four parameters, I can fit an elephant, and with five I can make him wiggle his trunk”.

Exponential regression

Power dependencies

Power dependencies