

Direction fields,
autonomous ODEs, and
the phase line

Lecture 7d: 2023-03-02

MAT A35 – Winter 2023 – UTSC

Prof. Yun William Yu


1st-order ODEs and slopes of solutions

- We can write a 1st-order ODE as $y' = f(x, y)$
- Recall that the derivative can be thought of as the slope of a solution.

Ex. $y' = x + y$


A sol passing through pt $(1, 1)$ has slope $y' = 2$

At pt $(0, 1)$, slope = 1



Direction field

- A direction field graphs out the slopes of all solutions going through a point.
- We can visualize different solutions by drawing trajectory curves that are always tangent to the direction field.



<https://www.wolframalpha.com/input/?i=slope+field+of+y%27%3Dx%2By>

Autonomous ODEs

- Recall that an autonomous ODE is one that does not have an explicit dependence on the independent variable (e.g. time).
- A first-order autonomous ODE can be rewritten in the form:

$$y' = f(y)$$

Ex. $y'' + y' + y = 0$ - 2nd order autonomous

Ex. $(y')^2 - \sin y - 1 = 0$ - 1st order autonomous


$$\Rightarrow (y')^2 = \sin y + 1$$

$$y' = \sqrt{\sin y + 1} \quad \text{standard form}$$


Direction fields of autonomous ODEs

- Notice that if $y' = f(y)$, then the slope has no x -dependence.

$$y' = \sin y$$



$$y' = y^2$$



Equilibrium values

- An equilibrium value of the autonomous ODE $y' = f(y)$ is a constant solution $y = c$.
- We can solve for equilibrium values by setting $y' = 0$.


Ex. $y' = y^2 = 0$
 $\Rightarrow y = 0$ is the only eq. value

Ex. $y' = \sin y = 0$
 $\Rightarrow y = k\pi$ for any integer k
are all equilibria.

Try it out: find the equilibrium values

• $y' = y^2 - y - 6$

$$y^2 - y - 6 = 0$$
$$(y - 3)(y + 2) = 0$$
$$y = 3, -2$$



unstable





asymptotically stable

- A: -2
- B: 3
- C: All of the above
- D: ???
- E: None of the above

Stability of equilibrium values

• Consider an equilibrium value c of $y' = f(y)$, and an initial value $y(0) = y_0$, where $y_0 \approx c$, but $y_0 \neq c$. Then c is

- Unstable if y diverges from c as time $x \rightarrow \infty$
- Asymptotically stable if $y(x) \rightarrow c$ as $x \rightarrow \infty$
- Semi-stable if as $x \rightarrow \infty$, $y(x)$ goes to c on one side, but diverges on the other side.



Determining stability using sign of y'

- $y' = 0$ at equilibrium.
- $y' > 0$ implies $y(x)$ gets bigger
- $y' < 0$ implies $y(x)$ gets smaller



Ex. $y' = y^2 - y - 6$
 $y' = (y-3)(y+2)$

unstable

If $y < -2$, $y' = (-)(-) = (+) > 0$


$-2 < y < 3$, $y' = (-)(+) = (-) < 0$

$3 < y$, $y' = (+)(+) > 0$







Phase line

- Draw arrows along the y-axis depending on if $z = f(y) = y'$ is positive or negative.



Phase line stability of $y' = f(y)$


- If $z = f(y)$ crosses the y -axis at $f(c) = 0$ going upward, then c is unstable.
- If $z = f(y)$ crosses the y -axis at $f(c) = 0$ going downward, then c is asymptotically stable.
- If $z = f(y)$ touches the y -axis at $f(c) = 0$, but remains on the same side of the y -axis, then c is semi-stable.



Try it out

$$\frac{dy}{dx} = f(y)$$


- $y' = y^5 - 3y^4 - 4y^3 + 12y^2$
- Classify the stability of the following equilibria:
- $y = -2$ *unstable*
- $y = 0$ *semi*
- $y = 2$ *stable*
- $y = 3$ *unstable*



- A: Asymptotically stable
- B: Unstable
- C: Semi-stable
- D: ???
- E: None of the above

Derivative test for stability

- Let $y' = f(y)$ have an equilibrium at $f(c) = 0$.
- If $\frac{df}{dy}(c) > 0$, then $y = c$ is unstable.
- If $\frac{df}{dy}(c) < 0$, then $y = c$ is asymptotically stable.
- If $\frac{df}{dy}(c) = 0$ and c is a local extremum (max or min) of $f(y)$, then $y = c$ is semi-stable



Example

$$y' = \frac{dy}{dx}$$

$$y' = 0$$

What is y ?

$$y' = f(y)$$

• $y' = y^5 - 3y^4 - 4y^3 + 12y^2$ has equilibria -2, 0, 2, 3

$$f(y) = y^5 - 3y^4 - 4y^3 + 12y^2$$

$$\frac{df}{dy} = 5y^4 - 12y^3 - 12y^2 + 24y$$

$$\frac{df}{dy}(-2) = 80 + 96 - 48 - 48 = 80 > 0, \quad \text{so } -2 \text{ is unstable}$$

$$\frac{df}{dy}(0) = 0$$


$$\frac{d^2f}{dy^2} = 20y^3 - 36y^2 - 24y + 24$$

$$\frac{d^2f}{dy^2}(0) = 24, \quad \text{so } 0 \text{ is a local min of } f(y),$$

$\Rightarrow 0 \text{ is semistable}$

$$\frac{df}{dy}(2) = 80 - 96 - 48 + 48 = -16, \quad \text{so } 2 \text{ is stable}$$

$$\frac{df}{dy}(3) = 45, \quad \text{so } 3 \text{ is unstable}$$



Try it out



https://commons.wikimedia.org/wiki/File:Glass_of_milk.jpg

- Fat crystallization: Let $y(x)$ be the proportion of crystallizable milk fat in a sample after x hours, satisfying

$$\frac{dy}{dx} = y' = 8(y^5 - y)$$

- If you start with half of the fat as crystallizable, how much fat is crystallizable as time goes to ∞ ?

- A: All of the fat
- B: Half of the fat
- C: None of the fat
- D: ???
- E: None of the above


$$y' = 0 = 8(y^5 - y) = 8y(y^4 - 1) = 8y(y^2 + 1)(y^2 - 1)$$

$$0 = y(y^4 - 1)$$


$$0 = y(y^2 + 1)(y^2 - 1)$$

$$0 = y(y^2 + 1)(y + 1)(y - 1)$$

$$y = -1, 0, 1$$



Logistic Growth Model



- Previously, we saw exponential growth $y' = ky$, $y(0) = y_0$, which had a solution $y(x) = y_0 e^{kx}$.
- In practice, this is unrealistic. For example, bacteria in a petri dish will initially grow almost exponentially, but then they'll use up all the available media.
- A better model is the logistic model, $y' = ky \left(1 - \frac{y}{L}\right)$, where the parameter L is the carrying capacity of the environment, and $k > 0$ is still the growth rate.

Stability of equilibria of logistic model

$$\bullet y' = \underline{ky \left(1 - \frac{y}{L}\right)}$$


Solve for equilibria.

$$y' = 0 = ky \left(1 - \frac{y}{L}\right)$$

$$y = 0 \quad \text{or} \quad 1 - \frac{y}{L} = 0$$

$$\Rightarrow y = L$$

Two equilibria: 0, L



0 is unstable

L is stable

Logistic model behavior

- $y' = ky \left(1 - \frac{y}{L}\right)$
- $y < 0$ is not physical, since we cannot have negative bacteria.
- When $0 < y < L$, $y' > 0$, so the number of bacteria increase, up to L .
- When $y > L$, $y' < 0$, so the number of bacteria decrease, down to L .
- On large time-scales, we therefore have L bacteria.

